

# RSV & SCDM

## Shared Clinical Decision-Making

*Shared clinical decision-making (SCDM) occurs when a health care provider and a patient work together to make a health care decision that is best for the patient. The optimal decision takes into account evidence-based information about available options, the provider's knowledge and experience, and the patient's values and preferences.*

### RSV Vaccine Updates

In June 2023, ACIP recommended that adults aged 60 years and older may receive a single dose of RSV vaccine using SCDM.

As part of this discussion, providers and patients should consider the patient's risk for severe RSV-associated disease. Epidemiologic evidence indicates that persons aged 60 years and older who are at highest risk for severe RSV disease include those with:

- ✓ Chronic medical conditions such as lung, heart, kidney, or liver disease
- ✓ Moderate to severe immune compromise, due to a medical condition or treatment
- ✓ Residing in nursing homes, or other long-term care facilities

### RSV Vaccine Clinical Guidance

Coadministration of RSV vaccines with other adult vaccines during the same visit is acceptable. Available data on immunogenicity and reactogenicity of RSV vaccines when administered with other vaccines is currently limited.



### The **SHARE** Approach

**S**eek your patient's participation

**H**elp your patient explore and compare treatment options

**A**ssess your patient's values and preferences

**R**each a decision with your patient

**E**valuate your patient's decision